

SEPARATION THEOREMS FOR COMPACT HAUSDORFF FOLIATIONS

WOJCIECH KOZŁOWSKI AND SZYMON M. WALCZAK

ABSTRACT. We investigate compact Hausdorff foliations on compact Riemannian manifolds in the context of the Gromov-Hausdorff distance theory. We give some sufficient conditions for such foliations to be separated in the Gromov-Hausdorff topology (GH-separation theorem).

1. INTRODUCTION

The concept of the Gromov-Hausdorff distance (briefly GH-distance), being a generalization of the notion of the Hausdorff distance, was originally introduced by M. Gromov [3] in the late 1970s. Next, Gromov, Katsuda, Peters and others showed that the GH-distance theory applied to the Riemannian manifolds leads to remarkable results [1], e.g., the *Cheeger's finiteness theorem* follows from the *Gromov-Katsuda convergence theorem* [4].

In [7] and [8], the second named author investigated warped compact Hausdorff foliations from the GH-distance theory point of view. He gave a necessary and sufficient conditions for a sequence of warped compact Hausdorff foliation to be converged to the space of leaves with quotient metric.

In the light of the results there appears a natural question: *Suppose that two compact metric spaces X and X' are GH-close. Are always the compact Hausdorff foliations $(M, \mathcal{F}, g)$ and $(M', \mathcal{F}', g')$ with space of leaves coinciding with X and X' , respectively, GH-close?*

In this paper we show that for compact Hausdorff foliations the answer is negative (*GH-separation Theorem* - Theorem 3 - the main result of the paper).

2. GROMOV-HAUSDORFF DISTANCE

Let $C, K \subset X$ be compact subsets of a metric space (X, d) . The number

$$d_H(C, K) = \inf\{\epsilon > 0 : C \subset N(K, \epsilon) \wedge K \subset N(C, \epsilon)\},$$

where $N(A, \epsilon) = \{x \in X : d(x, A) < \epsilon\}$, is called the *Hausdorff distance* of C and K .

Let (X, d_X) and (Y, d_Y) be arbitrary compact metric spaces. Equip the disjoint union $X \amalg Y$ with an admissible metric d , i.e., the metric which extends d_X and d_Y . The *Gromov-Hausdorff distance* (cf. [1], [3] and [6]) (GH-distance) d_{GH} of the spaces (X, d_X) and (Y, d_Y) one can define as

$$d_{\text{GH}}(X, Y) := \inf\{d_{\text{H}}(X, Y)\},$$

where the infimum is taken over all admissible metrics on $X \amalg Y$. Note that two compact metric spaces are isometric iff their GH-distance equals zero. Consequently, GH-distance in the class of all classes of isometry of compact metric spaces with the GH-distance is a metric.

Lemma 1. *If there exist ϵ -nets $\{x_1, \dots, x_k\} \subset X$ and $\{y_1, \dots, y_k\} \subset Y$ satisfying for all $1 \leq i, j \leq k$*

$$|d_X(x_i, x_j) - d_Y(y_i, y_j)| \leq \epsilon$$

then $d_{\text{GH}}(X, Y) \leq 3\epsilon$.

Proof. For a proof we refer to [1]. □

Lemma 2. *If $d_{\text{GH}}(X, Y) \leq \epsilon$ then for every ϵ -net $\{x_1, \dots, x_k\} \subset X$ there exists a 3ϵ -net $\{y_1, \dots, y_k\} \subset Y$ such that $|d_X(x_i, x_j) - d_Y(y_i, y_j)| \leq 2\epsilon$ for all $1 \leq i, j \leq k$.*

Proof. Since $d_{\text{GH}}(X, Y) \leq \epsilon$ there exists an admissible metric d on $X \amalg Y$ such that the Hausdorff distance $d_{\text{GH}}(X, Y) \leq d_{\text{H}}(X, Y) \leq \epsilon$. Let $\{x_1, \dots, x_k\}$ be an ϵ -net on X . For every $i \in \{1, \dots, k\}$ there exists $y_i \in Y$, $y_i \neq y_j$ while $i \neq j$, such that $d(x_i, y_i) \leq \epsilon$. Since d is an extension of metrics d_X and d_Y , $\{y_1, \dots, y_k\}$ is a 3ϵ -net on Y . Moreover,

$$d_X(x_i, x_j) \leq d(x_i, y_i) + d(y_i, y_j) + d(y_j, x_j) \leq d_Y(y_i, y_j) + 2\epsilon,$$

and similarly $d_Y(y_i, y_j) \leq d_X(x_i, x_j) + 2\epsilon$. □

Let $\text{Cov}_\epsilon(X)$ denote the smallest number of open ϵ -balls which covers X , and $\text{Cap}_\epsilon(X)$ the largest number of disjoint ϵ -balls contained in X . Obviously

$$\text{Cap}_\epsilon(X) \leq \text{Cov}_\epsilon(X).$$

Lemma 3. *Let (X, d) be a compact metric space. Then $\text{Cov}_{2\delta}(M) \leq \text{Cap}_\delta(M)$. More precisely, if $x_1, \dots, x_{\text{Cap}_\delta(M)}$ are the centres of disjoint balls $B(x_i, r)$ then the balls $B(x_i, 2r)$ cover M .*

Proof. Let $k = \text{Cap}_\delta(X)$ and $B(x_1, \delta), \dots, B(x_k, \delta)$ be a family of open disjoint balls in X . Let $x \in X$. Then $B(x, \delta) \cap B(x_i, \delta) \neq \emptyset$, and $d(x, y) < \delta$ and $d(y, x_i) < \delta$. Therefore, $x \in B(x_i, 2\delta)$ for some $i \in \{1, \dots, k\}$. Hence, $B(x_1, 2\delta), \dots, B(x_k, 2\delta)$ cover X . □

Lemma 4. *If $\text{Cov}_\epsilon(X) < \text{Cap}_{3\epsilon}(Y)$ then $d_{\text{GH}}(X, Y) > \epsilon$.*

Proof. Suppose that $d_{\text{GH}}(X, Y) \leq \epsilon$. Let $\{x_1, \dots, x_k\}$, $k = \text{Cov}_\epsilon(X)$, be an ϵ -net on X . Then, by Lemma 2, there exists a 3ϵ -net $\{y_1, \dots, y_k\}$ on Y , so $\text{Cov}_{3\epsilon}(Y) \leq \text{Cov}_\epsilon(X)$. Thus

$$\text{Cov}_{3\epsilon}(Y) \leq \text{Cov}_\epsilon(X) < \text{Cap}_{3\epsilon}(Y) \leq \text{Cov}_{3\epsilon}(Y).$$

Contradiction gives us the statement. $\square$

Let $p \geq 0$. A Borel measure μ on a metric space (X, d) is called a *p-dimensional Bishop measure* on (X, d) if there exist constants $\beta \geq 1$ and $\eta_0 > 0$ such that for all $\eta < \eta_0$ and every $x \in X$

$$(1) \quad \frac{1}{\beta} \eta^p \leq \mu(B(x, \eta)) \leq \beta \eta^p,$$

where $B(x, \eta) = \{y \in X : d(x, y) < \eta\}$.

Let (X, d) be a length-space, i.e. $d(x, y) = \inf\{l(\gamma)\}$, where $\gamma : [0, 1] \rightarrow X$ is a curve such that $\gamma(0) = x$, $\gamma(1) = y$, and $l(\gamma)$ denotes the length of γ .

Lemma 5. *If the balls $B(x, \delta)$ and $B(y, \delta)$ are disjoint, then $d(x, y) \geq 2\delta$.*

Proof. Suppose that $d(x, y) < 2\delta$. Let $\gamma : [0, 1] \rightarrow X$ be a curve from $x = \gamma(0)$ to $y = \gamma(1)$ with its length $l(\gamma) < 2\delta$. Let $t_0 \in [0, 1]$ be such that

$$l(\gamma|[0, t_0]) = l(\gamma|[t_0, 1]) = \frac{1}{2} l(\gamma) \leq \delta.$$

If $z = \gamma(1/2)$ then $d(x, z) < \delta$ and $d(z, y) < \delta$. Thus, $z \in B(x, \delta) \cap B(y, \delta)$. Contradiction ends our proof. $\square$

Lemma 6. *Let (X, d) be a compact length space, $p \geq 1$, and let μ be a p-dimensional Bishop measure on (X, d) with constants $\beta > 1$, $\eta_0 > 0$. There exist positive constants $C \geq 1$ and $\theta > 0$ such that for every $0 < r < \theta$ and $x \in M$,*

$$\frac{1}{Cr^p} \mu(X) \leq \text{Cap}_r(X) \leq \frac{C}{r^p} \mu(X).$$

Proof. Let $0 < r < \eta_0$, $k = \text{Cap}_r(X)$, and let $B(x_1, \delta), \dots, B(x_k, \delta)$ be a family of open disjoint balls in X . By (1), $\mu(B(x_i, r)) \geq \beta^{-1} r^p$, and

$$(2) \quad \mu(X) \geq \sum_{i=1}^k \mu(B(x_i, r)) \geq k \cdot \frac{1}{\beta} r^p.$$

Let $0 < r < \eta_0/2$. By Lemma 3, we have

$$(3) \quad \mu(X) \leq \sum_{i=1}^k \mu(B(x_i, 2r)) \leq k \cdot \beta (2r)^p.$$

Putting $C = \beta 2^p$ and $\theta = \eta_0/2$, (2) and (3) give us the statement. $\square$

Corollary 1. *Let $0 < r < \theta$ and $\alpha > 0$ be such that $r\alpha < \theta$. Then*

$$\alpha^{-p}C^{-2}\text{Cap}_r(X) \leq \text{Cap}_{\alpha r}(X) \leq \alpha^{-p}C^2\text{Cap}_r(X).$$

Remark 1. *Note that the volume form on a n -dimensional compact Riemannian manifold defines an n -dimensional Bishop measure.*

3. COMPACT HAUSDORFF FOLIATIONS

A foliation with all leaves compact is called a *compact foliation*. Let us consider any compact foliation $\mathcal{F}$ on a manifold M , and let $\pi : M \rightarrow \mathcal{L}$ denote a quotient map onto the space of leaves $\mathcal{L}$, this means that π identifies each leaf to a point. The space of leaves often is non-Hausdorff. Due to the results by D.B.A. Epstein [2], we recall theorems that describe the topology of such foliation:

Theorem 1. *The following conditions are equivalent.*

- (i) π is a closed map.
- (ii) π maps compact sets onto closed sets.
- (iii) Each leaf has arbitrarily small saturated neighbourhoods.
- (iv) $\mathcal{L}$ with quotient topology is Hausdorff.
- (v) If $K \subseteq M$ is compact, then the saturation of K is also compact.

Proof. For a proof we refer to [2], Theorem 4.1. □

Let M be a Riemannian manifold and N a submanifold on M . One can consider the induced Riemannian structure on N and introduce a volume of N as it's volume vol_N in the induced Riemannian structure. The next theorem describes the relation between the volume of the leaves defined above (briefly the volume function), the holonomy group of a leaf, and the topology of the space of leaves of a foliation $\mathcal{F}$ on a Riemannian manifold (M, g) .

Theorem 2. *If $(M, \mathcal{F}, g)$ is a foliated Riemannian manifold and L is a compact leaf of $\mathcal{F}$, then the following conditions are equivalent.*

- (i) *There exists a saturated neighbourhood N of the leaf L such that the volume function is bounded on N .*
- (ii) *The holonomy group of L is finite.*

Proof. For a proof we refer to [2], [Theorem 4.2]. □

The conditions of Theorem 2 imply that some saturated neighbourhood U of a compact leaf L consists of compact leaves, and in U the conditions of Theorem 1 are satisfied in U .

Moreover, by Reeb Stability Theorem, on a foliated manifold the conditions of Theorem 1 imply the conditions of Theorem 2.

A compact foliation which space of leaves is Hausdorff is called *compact Hausdorff foliation*. As an easy corollary of the above theorems we have:

Corollary 2. *Let $(M, \mathcal{F}, g)$ be a compact Riemannian manifold carrying compact Hausdorff foliation. Then $\sup_{L \in \mathcal{F}} \text{vol}(L) < \infty$.*

Now, let us consider the space of leaves $\mathcal{L}$ of an arbitrary compact Hausdorff foliation on a compact Riemannian manifold. Let us introduce on $\mathcal{L}$ a metric ρ defined by

$$\rho(L, L') = \inf \left\{ \sum_{i=1}^{n-1} \text{dist}(L_i, L_{i+1}) \right\},$$

where $L_1 = L$, $L_n = L'$, and the infimum is taken over all finite sequences of leaves. One can see that for a compact Riemannian foliation $\mathcal{F}$ the distance ρ coincides with Hausdorff distance of leaves of $\mathcal{F}$.

Remark 2. *Let g, g' be two Riemannian metrics on a compact foliated manifold $(M, \mathcal{F})$, where $\mathcal{F}$ is a compact Hausdorff foliation. Denote by ρ and ρ' two metrics on the space of leaves constructed using g and g' , respectively. Since M is compact, then $\frac{1}{C}g \leq g' \leq Cg$ for some constant $C \geq 1$. One can check that*

$$\frac{1}{C}\rho \leq \rho' \leq C\rho.$$

In further considerations we will need the following:

Lemma 7. *For every compact Hausdorff foliation $\mathcal{F}$ on a compact Riemannian manifold (M, g) there exists Riemannian structure $\tilde{g}$ on M such that $\mathcal{F}$ becomes a Riemannian foliation, and for any leaf $L \in \mathcal{F}$ we have $\tilde{g}|_L = g|_L$*

Proof. Obvious. See [5]. □

4. SEPARATION THEOREM

We say that a compact metric space (X', d') is *broadier* than a metric space (X, d) , and we briefly write $X' \succeq X$, if $\text{Cap}_\delta(X') \geq \text{Cap}_\delta(X)$ for all $\delta > 0$.

Let $d > 0$ be a real number. Let us denote by $\mathcal{M}(d, C, p, n)$ the class of all n -dimensional compact foliated Riemannian manifolds $(M, \mathcal{F}, g)$ carrying a compact Hausdorff foliation of dimension p satisfying:

- (1) For any leaf $L \in \mathcal{F}$, $\epsilon < d$, and any two balls $B_L(x, \epsilon), B_L(y, \epsilon)$ that are disjoint in L , the balls $B(x, \epsilon)$ and $B(y, \epsilon)$ are disjoint in M ;

- (2) $\max_{L \in \mathcal{F}} C_L \leq C$, $\min_{L \in \mathcal{F}} \theta_L \leq C$, $C_M \leq C$, $\theta_M \leq C$, where $C_L, \theta_L, C_M, \theta_M$ are the constants mentioned in Lemma 6 for a leaf L of $\mathcal{F}$ and for the manifold M , respectively;
- (3) $\frac{1}{C} \leq \text{vol}(L) \leq C$ for all $L \in \mathcal{F}$;
- (4) There exists a Riemannian structure $\tilde{g}$ on M satisfying $\frac{1}{C}g \leq \tilde{g} \leq Cg$ such that on $(M, \tilde{g})$ the foliation $\mathcal{F}$ becomes a compact Riemannian foliation.

Let $d > 0$, $C \geq 1$, and let $p, p', n, n' \in \mathbb{N}$ be such that $p' > p$ and $n' \geq n$.

Theorem 3. [GH-separation Theorem] *There exists $\epsilon > 0$ such that for any $(M, \mathcal{F}, g) \in \mathcal{M}(d_0, C, p, n)$ and $(M', \mathcal{F}', g') \in \mathcal{M}(d_0, C, p', n')$ such that $(M'/\mathcal{F}', \rho') \succeq (M/\mathcal{F}, \rho)$ we have $d_{\text{GH}}(M, M') > \epsilon$.*

Proof. Let $\mathcal{L} = M/\mathcal{F}$, $\mathcal{L}' = M'/\mathcal{F}'$, and let $\pi : M \rightarrow \mathcal{L}$ and $\pi' : M' \rightarrow \mathcal{L}'$ denote the natural projections. Let $r < \min\{d, C\}/3C$, and let $B(x_1, r/2), \dots, B(x_k, r/2)$, $k = \text{Cap}_{r/2}(\mathcal{L})$, be a family of open disjoint balls in $\mathcal{L}$. Since $\mathcal{L}' \succeq \mathcal{L}$ we can choose points $x'_1, \dots, x'_k$ in $\mathcal{L}'$ such that the balls $B(x'_1, r/2), \dots, B(x'_k, r/2)$ are also disjoint.

Now, in every leaf $L'_i = (\pi')^{-1}(x'_i)$ let us choose points

$$\xi'_{i,1}, \dots, \xi'_{i,l'},$$

where $l' = \min_{L \in \mathcal{F}'} \text{Cap}_{r/2}(L) > 0$, such that the balls $B_{L'_i}(x'_1, r/2), \dots, B_{L'_i}(x'_{l'}, r/2)$ are disjoint. Since $r < d$, the balls $B(\xi'_{i,j}, r/2)$ are disjoint in M' . Consequently, by Lemma 6, we have

$$(4) \quad \text{Cap}_{\frac{r}{2}}(M') \geq k \cdot l' \geq k \cdot \frac{2^{p'}}{\max_{L \in \mathcal{F}'} C_L r^{p'}} \min_{L \in \mathcal{F}'} \text{vol}'(L) \geq \frac{k}{C^2} \cdot \frac{2^{p'}}{r^{p'}},$$

where $\text{vol}'(L)$ denote the volume of a leaf in the induced Riemannian structure.

Now, let $\tilde{g}$ be a Riemannian structure on M mentioned in Lemma 7 such that $\mathcal{F}$ becomes a Riemannian foliation and such that

$$(5) \quad \frac{1}{C}g \leq \tilde{g} \leq Cg.$$

Let us choose in $L_i = \pi^{-1}(x_i)$ points $\xi_{i,1}, \dots, \xi_{i,l_i}$, $l_i = \text{Cap}_{r/2}(L_i)$ such the balls $B_{L_i}(\xi_{i,j}, r/2)$ are pairwise disjoint on $(L_i, \tilde{g}|_{L_i})$. By (5) and Remark 2, the balls $B(\xi_{i,j}, Cr)$ covers (M, g) , and

$$\text{Cov}_{Cr}(M) \leq k \cdot l,$$

where $l = \max_{i \in \{1, \dots, k\}} \{l_i\}$. Moreover, by (5) and Lemma 6,

$$l \leq \max_{L \in \mathcal{F}} \text{Cap}_{\frac{r}{2C}}(L) \leq \max_{L \in \mathcal{F}} (C_L) \frac{(2C)^p}{r^p} \cdot \max_{L \in \mathcal{F}} \text{vol}(L) \leq \frac{2^p C^{p+2}}{r^p},$$

and

$$\text{Cov}_{Cr}(M) \leq k \cdot \frac{2^p C^{p+2}}{r^p} =: B(r).$$

By Corollary 1,

$$\text{Cap}_{3Cr}(M') \geq \frac{1}{(6C)^{n'} \cdot (C)^2} \text{Cap}_{\frac{\pi}{2}}(M').$$

Next, by (4),

$$\text{Cap}_{3Cr}(M') \geq \frac{k}{(6C)^{n'} \cdot C^4} \cdot \frac{2^{p'}}{r^{p'}} =: A(r)$$

It follows that

$$\frac{A(r)}{B(r)} = \beta \cdot r^{p-p'}.$$

where β depends only on C , p' , and n' . Since $p < p'$ then $\lim_{r \rightarrow 0} \frac{A(r)}{B(r)} = +\infty$. Hence, there exists $\epsilon_0 < r$ such that $\text{Cap}_{3C\epsilon_0}(M') \geq A(r) > B(r) \geq \text{Cov}_{C\epsilon_0}(M)$. By Lemma 2, we obtain $d_{\text{GH}}(M, M') > C\epsilon_0$. $\square$

REFERENCES

- [1] D.-P. Chi and G. Yun, Gromov-Hausdorff Topology and its Applications to Riemannian Manifolds, Seoul National University, (1998).
- [2] D.B.A. Epstein, Foliations with all leaves compact, Ann. Inst. Fourier Grenoble **26** (1976), 265-282.
- [3] M. Gromov, Metric structures for Riemannian and non-Riemannian spaces, Birkhäuser, (1999).
- [4] A. Katsuda, Gromov's convergence theorem and its application, Nagoya Math. J. Vol. **100** (1985), pp. 11-48.
- [5] I. Moerdijk and J. Mrcun, Introduction to Foliations and Lie Groupoids, Cambridge University Press, (2003).
- [6] P. Petersen, Riemannian Geometry, Springer, (1997).
- [7] Sz. M. Walczak, Collapse of warped foliations, Diff. Geom. App., 25/6 (2007), 649-654.
- [8] Sz. M. Walczak, Warped compact foliations, Preprint of Faculty of Mathematics, University of Łódź, 2006/07.